

XXXIII INTERNATIONAL CONFERENCE ON HIGH ENERGY PHYSICS ICHEP'06

Moscow,
July 26-August 2, 2006

Organization

- Russian Academy of Sciences (RAS)
- The Ministry of Education and Science of RF
- Federal Agency of Science and Innovations of RF
- Federal Atomic Energy Agency of RF
- Joint Institute for Nuclear Research (JINR)
- Moscow State University (MSU)

Highlights from the Summer Conferences

Agnieszka Zalewska

SPC, 17.10.2006

Observation of B_s mixing at Tevatron was the most important experimental result presented at I CHEP06

but

Many other interesting results as well

- Around LHC
- Electroweak physics
- Neutrino physics
- Astrophysics and cosmology
- CKM matrix and CP violation
- Spectroscopy
- Heavy ions
- News from theory

Around LHC

LHC for itself

Prospects for 2007 and 2008

Tevatron vs LHC

Analysis methods developed for LHC are
successfully applied at Tevatron

Who will find SM Higgs?

LHC for ILC


Supersymmetry at LHC

HERA for LHC

PDF's from HERA are essential for LHC

2007

What data samples in 2007 ?




- Start to commission triggers and detectors with collision data (minimum bias, jets, ...) in real LHC environment
- Maybe first physics measurements (minimum-bias, underlying event, QCD jets, ...)?
- Observe a few $W \rightarrow l\nu$, $\Upsilon \rightarrow \mu\mu$, $J/\psi \rightarrow \mu\mu$?

F.Gianotti

Physics in 2007?


Example 2: Measurement of the underlying event in di-jet production



Particle multiplicity of underlying event obtained from the region transverse to the leading jet.

Comparison of plateau's between LHC and Tevatron will tell if detector performance, reconstruction tools and physics are under control

Multiplicity of charged particles with $p_T > 0.5$ GeV and $|\eta| < 1$ in region transverse to leading jet



- 15 days of data taking in 2007 enough to cover up to $p_T(\text{leading jet}) \sim 40$ GeV

Expected data

2008

With the first physics run in 2008 ($\sqrt{s} = 14 \text{ TeV}$)

1 fb⁻¹ (100 pb⁻¹) \equiv 6 months (few days) at $L = 10^{32} \text{ cm}^{-2}\text{s}^{-1}$
with 50% data-taking efficiency
→ may collect a few fb⁻¹ per experiment by end 2008

Channels (<u>examples</u> ...)	Events to tape for 100 pb ⁻¹ (per expt: ATLAS, CMS)	Total statistics from some of previous Colliders
$W \rightarrow \mu \nu$	$\sim 10^6$	$\sim 10^4$ LEP, $\sim 10^6$ Tevatron
$Z \rightarrow \mu \mu$	$\sim 10^5$	$\sim 10^6$ LEP, $\sim 10^5$ Tevatron
$t\bar{t} \rightarrow W b W b \rightarrow \mu \nu + X$	$\sim 10^4$	$\sim 10^4$ Tevatron
QCD jets $p_T > 1 \text{ TeV}$	$> 10^3$	---
$gg \quad m = 1 \text{ TeV}$	~ 50	---

With these data:

- Understand and calibrate detectors in situ using well-known physics samples
e.g. - $Z \rightarrow ee, \mu\mu$ tracker, ECAL, Muon chambers calibration and alignment, etc.
- $t\bar{t} \rightarrow b\bar{t} bjj$ jet scale from $W \rightarrow jj$, b-tag performance, etc.
- Measure SM physics at $\sqrt{s} = 14 \text{ TeV}$: W, Z, $t\bar{t}$, QCD jets ...
(also because omnipresent backgrounds to New Physics)


→ prepare the road to discovery it will take time ...

SM Higgs at LHC

2008?



Needed $\int L dt$ (fb^{-1}) per experiment

What about the SM Higgs boson ?



here discovery easier with gold-plated $H \rightarrow ZZ \rightarrow 4l$
 → by end 2008 ?

$H \rightarrow 4l$: narrow mass peak, small background
 $H \rightarrow WW \rightarrow l\nu l\nu$ (dominant at the Tevatron): counting channel (no mass peak)



F. Gianotti

SM Higgs at LHC




Later than 2008

Light Higgs : more difficult

$m_H \sim 115 \text{ GeV}$ 10 fb^{-1} : $S/\sqrt{B} \approx 4$ ATLAS

K-factors $\equiv \sigma(\text{NLO})/\sigma(\text{LO}) \approx 2$
for $H \rightarrow \gamma\gamma$ NOT included (conservative)

3 (complementary) channels with similar (small) significances:



- different production and decay modes
- different backgrounds
- different detector/performance requirements:
 - ECAL crucial for $H \rightarrow \gamma\gamma$ (in particular response uniformity) : $\sigma/m \sim 1\%$ needed
 - b-tagging crucial for $t\bar{t}H$: 4 b-tagged jets needed to reduce combinatorics
 - efficient jet reconstruction over $|\eta| < 5$ crucial for $q\bar{q}H \rightarrow q\bar{q}\tau\tau$:
forward jet tag and central jet veto needed against background

All three channels require very good understanding of detector performance and background control to 1-10% \rightarrow convincing evidence likely to come later than 2008 ...

Note: $WH \rightarrow l\nu b\bar{b}$ (dominant at the Tevatron) provides less sensitivity than $t\bar{t}H$ at LHC

Tevatron for LHC

- The LHC will inherit
 - Precise determination of Δm_s and constraints on CP phase in B_s sector β_{B_s}
 - Precision M_t ($\delta M_t = \pm 1.0-1.5 \text{ GeV}/c^2$) and M_w ($\delta M_w = 15-25 \text{ MeV}/c^2$)
 - A more restricted New Physics parameter space

– A higgs mass

From conclusions of D. Glenzinski

→ 95% CL exclusion for $m_H = 115-185 \text{ GeV}$ with 8 fb^{-1}

→ At 115 GeV
needs $\sim 3 \text{ fb}^{-1}$

→ At 160 GeV
needs $\sim 5 \text{ fb}^{-1}$



LHC for ILC

2008

Example of "early" discovery: Supersymmetry ?

If SUSY at TeV scale \rightarrow could be found "quickly" ... thanks to:


- large \tilde{q}, \tilde{g} cross-section $\rightarrow \approx 10$ events/day at 10^{32} for $m(\tilde{q}, \tilde{g}) \sim 1$ TeV
- spectacular signatures (many jets, leptons, missing E_T)



Our field, and planning for future facilities, will benefit a lot from quick determination of scale of New Physics. E.g. with 100 (good) pb^{-1} LHC could say if SUSY accessible to a ≤ 1 TeV ILC

BUT: understanding E_T^{miss} spectrum (and tails from instrumental effects) is one of the most crucial and difficult experimental issue for SUSY searches at hadron colliders.

HERA for LHC → beauty contribution to F_2




A.Zalewska, SPC, 17.10.2006

first measurement of $F_2^{b\bar{b}}$

first NNLO calculation

data in agreement with NLO
and NNLO


checks b PDF for LHC:



Electroweak physics

- W mass from LEP2
- Higgs mass from the global EW fit
- New measurements of τ lepton mass


W mass at LEP



•Main area of recent work: Final State Interactions (FSI)

- Mainly effects $qqqq$ channel
 - “color reconnection”
 - Bose-Einstein correlations

Summer 2006 - LEP Preliminary



prob. fit: 11.1%

Difference between channels:
 $M_W(4q-2q) = -11.7 \pm 44.6 \text{ MeV}/c^2$

Top quark mass


- Measured at the Tevatron in modes:

- Dilepton $t\bar{t} \rightarrow \ell \nu \ell \nu b\bar{b}$
- Lepton + jets $t\bar{t} \rightarrow \ell \nu jj b\bar{b}$
- All jets $t\bar{t} \rightarrow jjjj b\bar{b}$

- Many new results since last year
- New preliminary Tevatron average:


$$M_{top} = 171.4 \pm 2.1 \text{ GeV}$$

See Doug Glenzinski's talk (following directly) for description of top mass measurements




Global EW fit

Fit Constraints in M_W - M_t plane – March 06




Fit Constraints in M_W - M_t plane – July 06



**t mass smaller, W mass smaller
→ Higgs mass smaller**

SM Higgs Constraints

- EW fits alone, without theory uncertainties: $M_H = 85^{+39}_{-28}$ GeV (68% CL)
- 95% one-sided CL including theory uncertainties (“blue band”):
 - $M_H < 166$ GeV (ignoring direct limit)
 - $M_H < 199$ GeV (including 114 GeV limit)
- Blue band uncertainties due to uncalculated higher order corrections, estimated by ZFITTER



τ Lepton mass



New preliminary measurement of the τ mass from lower energy e^+e^- experiments

- KEDR at VEPP-4M
- BELLE at KEK


$$M_\tau = 1776.71 \pm 0.13(\text{stat}) \pm 0.32(\text{syst}) \text{ MeV} \quad (\text{BELLE, preliminary})$$

$$M_\tau = 1776.80 \pm_{0.22}^{0.25}(\text{stat}) \pm 0.15(\text{syst}) \text{ MeV} \quad (\text{KEDR, preliminary})$$

$$M_\tau = 1776.99 \pm_{0.26}^{0.29} \text{ MeV} \quad (\text{PDG})$$



KEDR
 $\tau^+\tau^-$ CROSS
 section near
 threshold



Neutrino physics

- ➔ Oscillations – status
- ➔ θ_{13} in reactor exp.: Double Chooz and Daya Bay
- ➔ SuperK III
- ➔ Solar neutrinos: Borexino and KamLAND
- ➔ MINOS – first year of data taking
- ➔ Prospects for OPERA
- ➔ Update of DONUT

Still waiting for MiniBooNE!

Neutrino oscillations

results can be encoded in a Lorentz-invariant Lagrangian

$$L = L_{SM} + \delta L(m_\nu) + \delta L$$



1st evidence of physics beyond the SM after more than 30 years!

additional operators giving negligibly small contributions to ν propagation in present experiments

L_{SM} invariant under

local $SU(3) \otimes SU(2) \otimes U(1)$

global, non - anomalous

$$\frac{B}{3} - L_e, \quad \frac{B}{3} - L_\mu, \quad \frac{B}{3} - L_\tau$$

broken individually by $\delta L(m_\nu)$
possible exception: $(B - L)$


either $\dim(\delta L) \geq 6$ such as e.g.

$$\frac{c_1}{\Lambda^2} \psi \psi l l + \frac{c_2}{\Lambda^2} (lH)^+ \gamma^\mu \partial_\mu (lH) + \dots$$

or **new particles** in δL such as

$$\lambda \phi \nu \nu + \dots$$

new (pseudo)scalar



$L_0=180\text{km}$ is used for KamLAND

F. Feruglio

low-energy parameters in $\delta L(m_\nu)$

ν masses

[3 light active ν]

m_1, m_2, m_3


order

$$m_1 < m_2$$

$$\Delta m_{21}^2 < \left| \Delta m_{32}^2 \right|, \left| \Delta m_{31}^2 \right| \quad [\Delta m_{ij}^2 \equiv m_i^2 - m_j^2]$$

i.e. 1 and 2 are, by definition, the closest levels

two, still open,
possibilities:



Mixing matrix (analogous to V_{CKM})

$$U_{PMNS} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{i\delta} \\ -s_{12}c_{23} - c_{12}s_{13}s_{23}e^{-i\delta} & c_{12}c_{23} - s_{12}s_{13}s_{23}e^{-i\delta} & c_{13}s_{23} \\ -c_{12}s_{13}c_{23} + s_{12}s_{23}e^{-i\delta} & -s_{12}s_{13}c_{23} - c_{12}s_{23}e^{-i\delta} & c_{13}c_{23} \end{pmatrix} \times \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\alpha} & 0 \\ 0 & 0 & e^{i\beta} \end{pmatrix}}$$

$$c_{12} \equiv \cos \theta_{12}, \dots$$

F.Feruglio

- only if ν are Majorana
- drops in oscillations

from data

[2σ errors (95% C.L.)]

F.Feruglio


$$\Delta m_{sol}^2 \equiv \Delta m_{21}^2 = 7.92 (1 \pm 0.09) \times 10^{-5} \text{ eV}^2$$

$$\Delta m_{atm}^2 \equiv |\Delta m_{32}^2| = 2.4 (1_{-0.26}^{+0.21}) \times 10^{-3} \text{ eV}^2$$

[Fogli, Lisi, Marrone, Palazzo, 0506083]

sign [Δm_{32}^2] unknown δ_{CP} unknown

$m_i < 2 \text{ eV}$ 95% C.L. [Tritium β -decay]



next CMB satellite + weak grav. lensing + improved galaxy survey $\sum m_i < (0.02 \div 0.08) \text{ eV} (1\sigma)$ [2015?]

$$\sin^2 \theta_{12} = 0.314 (1_{-0.15}^{+0.18})$$

$$\sin^2 \theta_{23} = 0.44 (1_{-0.22}^{+0.41})$$

$$\sin^2 \theta_{13} = 0.9_{-0.9}^{+2.3} \times 10^{-2}$$

two lepton mixing angles are large

$$V_{us} \approx \lambda \quad V_{cb} \approx \lambda^2 \quad V_{ub} \approx \lambda^3$$

$$\lambda \approx 0.22$$

δ, α, β unknown

some parameters measured already quite precisely

$$\theta_{12} = (34.1_{-1.6}^{+1.7})^\circ [1\sigma]$$

$$\theta_{12} + \underbrace{\theta_C}_{\text{Cabibbo angle}} = (47.0_{-1.6}^{+1.7})^\circ$$


quark-lepton complementarity?

[Raidal 0404046]

Minakata, Smirnov 0405088]

Most of plausible range for U_{e3} explored in 10 yr from now

based on
F.Feruglio



$|U_{e3}| < 0.05$ would select a very narrow (not empty) subset of existing models

θ_{13} – future

similar conclusion by:
Barbieri, Hambye, Romanino 0302118
Ibarra, Ross 0307051
Chen, Mahanthappa 0305088
Lebed, Martin 0312219
Joshipura @ NOON 2004

θ_{13} proposals

Reactor experiments

	Country	Reactor Power (GW _{th})	Distance (m)	Depth (mwe)	Far Target mass (t)	Sensitivity (90% u. l.)	Expected start date	Funding
Double CHOOZ	France	8.7	280 1050	80 300	10.2	0.03	2008 f 2009 f+n	○
KASKA	Japan	24.3	350 1600	90 260	8	0.025	2009/2010	△
RENO	Korea	17.3	150 1500	230 675	20x2	0.02	2009/2010	△
Daya Bay	China	11.6 17.4 (2011~)	360 (500) 1750	260 910	20x4	0.01	2009 n+m 2010 full	○
Angra	Brazil	4.1	300 1500	250 2000	500	0.005	-	-

Double CHOOZ – almost fully founded, Daya Bay – soon founded, is Double CHOOZ competitive?

Double CHOOZ

V.Sinev on July 29



use existing far site
quickest
baseline and depth are not ideal




statistical error 2.8% \Rightarrow 0.5%
systematic error 2.7% \Rightarrow ~0.6%

K.I noue

Daya Bay **Layout of the experiment**

K. Inoue

Y.Wang on July 29



Near-Far detector schemes:

To cancel reactor-related errors

Residual error ~0.1%

Swap near-far detectors

To cancel detector-related errors.

Residual error ~0.2%

Detector deep undergrounds

To reduce backgrounds

B/S at near site: ~0.5%

B/S at far site: ~0.2%

Fast Measurement

DYB+Mid, 2008-2009

Sensitivity (1 year) ~0.03



Full Measurement

DYB+LA+Far, from 2010

Sensitivity (3 year) <0.01


total sys.error 0.06~0.36%

Baseline, depth are optimized.



Super-Kamiokande III



Kun 30592 Sub 1 Ev 1
06-07-11:09:46:56
Inner: 927 bits, 4680 pR
Outer: 169 bits, 932 pR (11-CLM)
Trigger ID: 8x0b
S wall: 2699.0 cm
Fully-Correlated mode



SK full restoration has been completed and water is full since July 11, 2006. Detector calibration is going on.

Charge (pe)

- >26.7
- 23.3-26.7
- 20.0-23.3
- 17.3-20.2
- 14.7-17.3
- 12.2-14.7
- 10.0-12.2
- 8.0-10.0
- 6.2- 8.0
- 4.7- 6.2
- 3.3- 4.7
- 2.2- 3.3
- 1.3- 2.2
- 0.7- 1.3
- 0.2- 0.7
- < 0.2




First event in SK-III

K.I noue


Solar neutrinos: pp chain vs CNO cycle...

...coming back to David's original idea

Energy spectrum for the $\nu_e e$ elastic scattering



present spectrum in KamLAND




K.I noue

Borexino is filled with water,
in September KamLAND has started cleaning its scintillator

BOREXINO sees neutrinos from CERN (August 2006) !



L.Oberauer



Hall-C Side view

Borexino

CTF

Opera





Hall-C Top view



NuMI - MINOS


Construction 1998-2004; operations 2005 - 2010..





New Results from MINOS

Geoff Pearce, Rutherford Lab




Data sample	Data	Expected (Unoscillated)	Data/MC
ν_μ (<30 GeV)	215	336.0 ± 14.4	0.64 ± 0.05
ν_μ (<10 GeV)	122	238.7 ± 10.7	0.51 ± 0.05
ν_μ (< 5 GeV)	76	168.4 ± 8.8	0.45 ± 0.06



Closing in on Δm_{32}^2

R.Rameika



Submitted to hep-ex archive:
hep-ex 0607088

$$\Delta m_{32}^2 = 0.00274^{+0.44}_{-0.26} \times 10^{-3} \text{ eV}^2/\text{c}^4$$
$$\sin^2(2\theta_{23}) > 0.87 \text{ (68\% C.L.)}$$



OPERA

R.Rameika

τ decay channel	Signal		Background
	$\Delta m^2 = 2.4 \times 10^{-3} \text{ eV}^2$	$\Delta m^2 = 3.0 \times 10^{-3} \text{ eV}^2$	
$\tau \rightarrow \mu$	3.6	5.6	0.23
$\tau \rightarrow e$	4.3	6.7	0.23
$\tau \rightarrow h$	3.8	5.9	0.32
$\tau \rightarrow 3h$	1.1	1.7	0.22
ALL	12.8	19.9	1.0

full mixing, 5 years run @ 4.5×10^{19} pot / year

Higher Δm^2 from Minos very good for OPERA



Update from DONUT

R.Rameika



12 April 2001

PHYSICS LETTERS B

Phys. Lett. B 504 (2001) 218–224

www.elsevier.nl/locate/nucphys

Observation of tau neutrino interactions

DONUT Collaboration

K. Kodama^a, N. Ushida^a, C. Andreopoulos^b, N. Saoulidou^b, G. Tzanakos^b, P. Yager^c,
 B. Baller^d, D. Boehnlein^d, W. Freeman^d, B. Lundberg^d, J. Morfin^d, R. Rameika^d,
 J.C. Yun^d, J.S. Song^e, C.S. Yoon^e, S.H. Chung^e, P. Berghaus^f, M. Kubantsev^f,
 N.W. Reay^f, R. Sidwell^f, N. Stanton^f, S. Yoshida^g, S. Aoki^h, T. Hara^h, J.T. Rhee^h,
 D. Ciampaⁱ, C. Ericksonⁱ, M. Grahamⁱ, K. Hellerⁱ, R. Rusackⁱ, R. Schwienhorstⁱ,
 J. Stelaff^j, J. Trammell^j, J. Wilcox^k, K. Hoshino^l, H. Jiko^l, M. Miyanishi^l,
 M. Komatsu^m, M. Nakamura^m, T. Nakano^m, K. Niwaⁿ, N. Nonakaⁿ, K. Okadaⁿ, O. Satoⁿ,
 T. Akdogan^o, V. Paolone^o, C. Rosenfeld^o, A. Kulik^{o,1}, T. Kafka^o, W. Oliver^o,
 T. Patzak^o, J. Schneps^o

^a Aichi University of Education, Kariya, Japan

^b University of Illinois, Urbana, IL 61801, USA

^c University of California, Davis, Davis, CA 95616, USA

^d Fermilab, Batavia, IL 60510, USA

^e Gyeongang University, Cheju, South Korea

^f Kansas State University, Manhattan, KS 66506, USA

^g Kobe University, Kobe, Japan

^h Kyushu University, Fukuoka, Japan

ⁱ University of Minnesota, Minneapolis, MN 55455, USA

^j Nagoya University, Nagoya 464-8602, Japan

^k University of Pittsburgh, Pittsburgh, PA 15260, USA

^l University of South Carolina, Columbia, SC 29208, USA

^m Toho University, Tokyo, Japan

ⁿ Toho University, Tokyo, Japan

^o Toho University, Tokyo, Japan

¹ Toho University, Tokyo, Japan

² Toho University, Tokyo, Japan

³ Toho University, Tokyo, Japan

⁴ Toho University, Tokyo, Japan

⁵ Toho University, Tokyo, Japan

⁶ Toho University, Tokyo, Japan

⁷ Toho University, Tokyo, Japan

⁸ Toho University, Tokyo, Japan

⁹ Toho University, Tokyo, Japan

¹⁰ Toho University, Tokyo, Japan

¹¹ Toho University, Tokyo, Japan

¹² Toho University, Tokyo, Japan

¹³ Toho University, Tokyo, Japan

¹⁴ Toho University, Tokyo, Japan

¹⁵ Toho University, Tokyo, Japan

¹⁶ Toho University, Tokyo, Japan

¹⁷ Toho University, Tokyo, Japan

¹⁸ Toho University, Tokyo, Japan

¹⁹ Toho University, Tokyo, Japan

²⁰ Toho University, Tokyo, Japan

²¹ Toho University, Tokyo, Japan

²² Toho University, Tokyo, Japan

²³ Toho University, Tokyo, Japan

²⁴ Toho University, Tokyo, Japan

²⁵ Toho University, Tokyo, Japan

²⁶ Toho University, Tokyo, Japan

²⁷ Toho University, Tokyo, Japan

²⁸ Toho University, Tokyo, Japan


²⁹ Toho University, Tokyo, Japan

³⁰ Toho University, Tokyo, Japan

³¹ Toho University, Tokyo, Japan

³² Toho University, Tokyo, Japan

³³ Toho University, Tokyo, Japan



First 4 events published
in Phys. Lett. B 504, 218(2001)

Abstract

The DONUT experiment has analyzed 203 neutrino interactions recorded in nuclear emulsion targets. A decay search has found evidence of four tau neutrino interactions with an estimated background of 0.34 events. This number is consistent with the Standard Model expectation. © 2001 Published by Elsevier Science B.V.

PACS: 14.90.Ln


Now: 10 ν_τ candidates, publication of final results
by end of 2006

Astrophysics and cosmology

- ➔ Updates of EHE Cosmic Rays at 10^{19-20} eV
- ➔ Searches for Dark Matter

Y.Takahashi

Air shower observations for EHE particles → particle astronomy





Ground-based arrays

- (1) Scintillator array
- (2) Fluorescence telescope array

Energy Spectrum of Cosmic Rays

- LHC will comprehensively profile SUSY DM regime $m < 7 \text{ TeV}/c^2$
- the mystery of the EHECR Origins remains and extends to $m > 300 \text{ TeV}/c^2$ regime

Y. Takahashi



10^9 eV

10^{20} eV

$p + 2.7K \rightarrow e^+e^- + p$ above LHC $5 \cdot 10^{17} \text{ eV}$





$p + 2.7K \rightarrow + \rightarrow n + +, p + 37$

Y. Takahashi, D. C. ... 10/2000



4 EHE
Air Shower
Experiments:

AGASA,
Hi-Res,
Yakutsuk,
Auger

→ **Dip**
appears
common to all




Data Adjusted by Hill-Schramm-Berezinsky dip



$\lambda_{\text{AGASA}} = 0.9, \lambda_{\text{Yakutsk}} = 0.75, \lambda_{\text{HiRes}} = 1.2$; Each $\chi^2/\text{d.o.f} = 1.0 - 1.2$, adjusted for dips.
 (Berezinsky et al. 2006 PRD in press)

Confusing: AGASA preliminary new spectrum with recent Corsika simulator (Teshima)


Y. Takahashi



- Energy shift to lower direction
 $\sim 10\%$ at 10^{19}eV
 $\sim 15\%$ at 10^{20}eV
- Above 10^{20}eV
 11 events \rightarrow 5~6 events
- Featureless spectrum very close to E^{-3}
- P-SIBYLL (above 10^{19}eV)
 $\beta = 2.95 \pm 0.08$
 $(\chi^2 / \text{NDF} = 8.5/11)$
- Fe-QGSJET (above 10^{19}eV)
 $\beta = 2.90 \pm 0.08$
 $(\chi^2 / \text{NDF} = 8.5/11)$

EUSO (Extreme Universe Space Observatory) on ISS JEM

Y. Takahashi



10^{19} eV Track volume has S/N ~ 70




A. Zalewska, SPC, 17-18 2006
 NUV Fluorescence 300 - 400 nm; Čerenkov, too

EUSO



EUSO ~ 300 x AGASA ~ 10-50 x Auger


EUSO (Instantaneous) ~3000-15000 x AGASA ~ 100-500 x Auger



Y.Takahashi

Dark Matter searches -complementary approaches

Hunt for dark matter in the dark needs a set of different weapons, in an integrated way




Indirect searches

gamma-rays etc. on satellite



Direct searches in underground labs



LHC experiments

neutrinos in ice



neutrinos deep underwater

gamma-rays on surface



Principles of WIMP Detection

Target = Detector

Measure the energy deposited by the hit nucleus


Part of this energy appears as **charge, light or heat**

Main challenges

- Signal rate is small
- Energy deposit is tiny (several keV)
- Signal spectrum decreases exponentially

•3 basic backgrounds

- electromagnetic (β & γ); dominant \Rightarrow electrons
- neutrons (and WIMPs) \Rightarrow nuclear recoil
- surface contamination (partial energy release)



Against backgrounds, in Underground laboratory

Search for characteristic signal: annual modulation (DAMA only)

Talk by P.L. Belli

Passive shielding. Against external backgrounds only

Active discrimination. Against external and internal background

Measure two quantities

Event by event rejection. Necessary for “zero” background

Statistical rejection not enough: background might be signal, in presence of residual

background sensitivity proportional to square-root of exposure

Liquid Argon. WARP @ LNGS

Fast/slow components of light pulse in Ar have very different lifetimes: 7 ns/1600 ns
⇒ second powerful handle for discrimination

But huge intrinsic background: ^{39}Ar β decay @ 1 Bq/kg. Needs 10^{-8} rejection @ 10^4 kg d

140 kg TPC in construction @ LNGS




First tests (and physics) with 2.3 l prototype

Exposure = 96.5 kg d ⇒

Rejct power for $30 < E < 100 \text{ keV}_{ee} = 37 \cdot 1 \cdot 10^{-6}$
0 events @ $E > 42 \text{ keV}_{ee}$ (5 events @ $30 < E < 42$)

Caveat: Quenching factor not yet measured



A. Bettini

CKM matrix and CP violation

- ➔ B factories – fantastic performance
- ➔ CP violation in B decays – new measurements
- ➔ $|V_{ud}|$ from neutron lifetime measurement

No sign of New Physics

- ➔ but DD mixing is now considered to be a good probe of NP
- ➔ Future K experiments


Two Asymmetric-energy *B* Factories

M.Hazumi


PEP-II at SLAC

9GeV (e^-) \times 3.1GeV (e^+)
 peak luminosity:

$$1.12 \times 10^{34} \text{cm}^{-2}\text{s}^{-1}$$




BaBar




11 nations,
 80 institutes,
 623 persons

10.2006

13 countries,
 57 institutes,
 ~400 collaborators



Belle



KEKB at KEK

8GeV (e^-) \times 3.5GeV (e^+)
 peak luminosity:


$$1.65 \times 10^{34} \text{cm}^{-2}\text{s}^{-1}$$

world record

Integrated Luminosity

World Integrated Luminosity (KEKB+PEP-II)

As of July 24, 2006



10000/fb !!


reached on July 13, 2006

~ 1 Billion $B\bar{B}$ pairs

- Far beyond the design luminosities of both proposals
- Triumph in accelerator science

Unitarity triangle

Various observable quantities in B physics constrain the angles and lengths of the unitarity triangle.



Angle determination

- ϕ_1/β : **CP asymmetries in $b \rightarrow ccs$ modes** – well measured
- ϕ_2/α : **CP asymmetries in $B \rightarrow \pi\pi, \rho\rho, \rho\pi$** . – worse measured
- ϕ_3/γ : **CP asymmetries in $B \rightarrow DK$, etc.** - practically unconstrained
- $(2\phi_1 + \gamma)/(2\beta + \gamma)$: **CP asymmetry in $B \rightarrow D\pi$, etc**

New measurements of ϕ_1

- ϕ_1 with tree diagram ($b \rightarrow ccs$)
 - **BaBar:** $\sin 2\beta$ in $B^0 \rightarrow J/\psi K^0, \psi(2S)K_S, \eta_c K_S, \chi_{c1} K_S, J/\psi K^{*0}$
 - **Belle:** $\sin 2\phi_1$ in $B^0 \rightarrow J/\psi K^0$
- ϕ_1 with penguin diagram ($b \rightarrow sqq$)
 - **BaBar:** $B^0 \rightarrow K^+ K^- K^0, \eta' K^0, K_S K_S K_S, \omega K_S, \rho K_S, \pi^0 K_S$
 - **Belle:** $B^0 \rightarrow \eta' K^0, \pi^0 K_S$
 - **Belle:** $B^0 \rightarrow \phi K^0, K^+ K^- K_S, K_S K_S K_S, f^0 K_S, \omega K_S$

M.Hazumi


ϕ_1

Time-dependent CP violation (tCPV)

“double-slit experiment” with particles and antiparticles


Quantum interference b/w two diagrams

tree diagram

 J/ψ K_S

+


box diagram + tree diagram

 V_{td} K_S J/ψ V_{td}

You need to “wait” (i.e. $\Delta t \neq 0$) to have the box diagram contribution.


$B^0 \rightarrow J/\psi K^0 : 532 \text{ M } BB \text{ pairs}$

$B^0 \rightarrow J/\psi K_S^0$



$$M_{bc} = \sqrt{E_{beam}^{*2} - P_{J/\psi K_S}^{*2}}$$


$B^0 \rightarrow J/\psi K_L^0$



p_{KL} information is poor
→ lower purity



$B^0 \rightarrow J/\psi K^0$: combined result



Preliminary

532 M $B\bar{B}$ pairs

previous measurement
 $\sin 2\phi_1 = 0.652 \pm 0.044$
(386 M $B\bar{B}$ pairs)

$\sin 2\phi_1 = 0.642 \pm 0.031$ (stat) ± 0.017 (syst)
 $A = 0.018 \pm 0.021$ (stat) ± 0.014 (syst)



BaBar2006: $\sin 2\beta$ in $b \rightarrow ccs$

$$\sin 2\beta = 0.710 \pm 0.034(\text{stat}) \pm 0.019(\text{syst})$$

$$|\lambda| = 0.932 \pm 0.026(\text{stat}) \pm 0.017(\text{syst})$$




Preliminary

BABAR:CONF-06/036




$$A = -0.07 \pm 0.028 \pm 0.018$$



$B^0 \rightarrow J/\psi K^0,$
 $\psi(2S)K_S,$
 $\eta_c K_S,$
 $\chi_{c1} K_S,$
 $J/\psi K^{*0}$

α/ϕ_2

ϕ_1 “beam”



ϕ_2 “banana”

$$B \rightarrow \pi^+ \pi^-, \pi^\pm \pi^0, \pi^0 \pi^0$$

$$B \rightarrow \rho^0 \rho^0, \rho^\pm \rho^0, \rho^+ \rho^-$$

$$B^0 \rightarrow (\rho\pi)^0$$



Belle 2006: $B^0 \rightarrow \pi^+ \pi^-$ decay (CP asymmetry)

H. Ishino

532M $B\bar{B}$

BELLE-CONF-0649

1464 ± 65 signal events

$\pi^+ \pi^-$ yields

$\pi^+ \pi^-$ asymmetry

$$A_{\pi\pi} = +0.55 \pm 0.08 \pm 0.05$$

$$S_{\pi\pi} = -0.61 \pm 0.10 \pm 0.04$$



first error: stat., second: syst.

Large Direct CP violation (5.5σ)

Large mixing-induced CP violation (5.6σ)



confidence level contour



A.Zalewska, SPC, 17.10.2006

background subtracted

2 body π -K combinations

	BaBar		Belle	
	$BR \times 10^{-6}$	A_{cp}	$BR \times 10^{-6}$	A_{cp}
$B^+ \rightarrow \pi^+ \pi^0$	$5.12 \pm 0.47 \pm 0.29$	$-0.019 \pm 0.088 \pm 0.014$	$6.6 \pm 0.4^{+0.4}_{-0.5}$	$+0.07 \pm 0.06 \pm 0.01$
$B^0 \rightarrow \pi^+ \pi^-$	$5.8 \pm 0.4 \pm 0.3$	$+0.16 \pm 0.11 \pm 0.03$	$5.1 \pm 0.2 \pm 0.2$	$+0.55 \pm 0.08 \pm 0.05$
$B^0 \rightarrow \pi^0 \pi^0$	$1.48 \pm 0.26 \pm 0.12$	$+0.33 \pm 0.36 \pm 0.08$	$1.1 \pm 0.3 \pm 0.1$	$+0.44^{+0.73+0.04}_{-0.62-0.06}$
$B^0 \rightarrow K^+ \pi^-$	$19.7 \pm 0.6 \pm 0.6$	$-0.108 \pm 0.024 \pm 0.007$	$20.0 \pm 0.4^{+0.9}_{-0.8}$	$-0.093 \pm 0.018 \pm 0.008$
$B^0 \rightarrow K^0 \pi^0$	$10.5 \pm 0.7 \pm 0.5$	$-0.20 \pm 0.16 \pm 0.03$	$9.2^{+0.7+0.6}_{-0.6-0.7}$	$-0.05 \pm 0.14 \pm 0.05$
$B^+ \rightarrow K^+ \pi^0$	$13.3 \pm 0.56 \pm 0.64$	$+0.016 \pm 0.041 \pm 0.010$	$12.4 \pm 0.5^{+0.7}_{-0.6}$	$+0.07 \pm 0.03 \pm 0.01$
$B^+ \rightarrow K^0 \pi^+$	$23.9 \pm 1.1 \pm 1.0$	$-0.029 \pm 0.039 \pm 0.010$	$22.9^{+0.8}_{-0.7} \pm 1.3$	$+0.03 \pm 0.03 \pm 0.01$
$B^0 \rightarrow K^0 \bar{K}^0$	$1.08 \pm 0.28 \pm 0.11$	$0.40 \pm 0.41 \pm 0.06$	$0.86^{+0.24}_{-0.21} \pm 0.09$	$-0.57^{+0.72}_{-0.65} \pm 0.13$
$B^0 \rightarrow K^+ K^-$	< 0.40		< 0.25	
$B^+ \rightarrow \bar{K}^0 K^+$	$1.61 \pm 0.44 \pm 0.09$	$0.10 \pm 0.26 \pm 0.03$	$1.22^{+0.33+0.13}_{-0.28-0.16}$	$+0.13^{+0.23}_{-0.24} \pm 0.02$




Belle 2006: ϕ_2 from $B \rightarrow \rho\pi$


M.Hazumi

Dalitz analysis + isospin (pentagon) analysis

- 26(Dalitz) + 5($\text{Br}(\rho^\pm\pi^\pm)$, $\text{Br}(\rho^+\pi^0)$, $\text{Br}(\rho^0\pi^+)$, $A(\rho^+\pi^0)$, and $A(\rho^0\pi^+)$)



Preliminary Results




$$\alpha/\phi_2 = [83^{+12}_{-23}]^\circ \quad (1\sigma)$$

(no constraint at 2σ)

will be included in the world average

A.Zalewska, SPC, 17.10.2006

D⁰-D⁰ mixing status



NO MIXING $(x,y)=(0,0)$ excluded:

- ✓ $\sim 2.1\sigma$ Belle $D^0 \rightarrow K\pi$ (assuming no CPV)
- ✓ $\sim 2.3\sigma$ BaBar $D^0 \rightarrow K2\pi/K3\pi$
- ✓ $\sim 2.2\sigma$ Average y_{CP}

Many results coming soon:

- BaBar $D^0 \rightarrow K\pi$
- Belle $D^0 \rightarrow K2\pi/K3\pi$
- BaBar & Belle updates of y_{CP}
(5 × data set available)
- BaBar semileptonic
- BaBar & Belle $D^0 \rightarrow K_S \pi\pi$ (Dalitz)

CLEOc with projected data set can reach UL on $R_M \sim \text{few} \times 10^{-4}$ (on $x \sim 2\%$)

D⁰D⁰ mixing observation may be on the horizon


Super B factory

Current Belle+BaBar integrated luminosity: $L(\text{now}) \sim 1/\text{ab}$

In many aspects asymmetric B factories are complementary to B physics in hadron machine (unique for neutrino and tau modes)


5X L(now)

50X L(now)



Neutron lifetime measurements $\rightarrow V_{ud}$

A.Srebrov




Neutron lifetime measurements with gravitational trap of ultracold neutrons

PNPI-ILL-JINR

Setup for the measurement
of n-lifetime at ILL
(Grenoble, France)



Neutron decay and Standard Model



$$A = -0.1189(8) \quad \text{PERKEO 2002}$$

$$\tau_n = 885.7 \pm 0.8 \text{ s} \quad \text{PDG(2003)}$$

$${}^n V_{ud} = 0.9717(13)$$

$${}^{00} V_{ud} = 0.9738(5)$$

$$V_{us} = 0.2196(23) \quad \text{PDG(2003)}$$

$$V_{ub} = 0.0036(9) \quad \text{PDG(2003)}$$

$$|{}^n V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1 - \Delta = 0.9924(28)$$

$$\Delta = 0.0076(28) = 2.7\sigma$$

$${}^n V_{ud} - {}^{00} V_{ud} = -0.0021(26) = -0.8\sigma$$

Neutron decay and cosmology

G. J. Mathews, T. Kajino, T. Shima, Phys. Rev. D 71, 021302(R) (2005)



$$(f\tau_n)^{-1} = \frac{G_F^2}{2\pi^3} (1 + 3g_A^2) m_e^5$$

$$\Gamma = (7/60)\pi(1 + 3g_A^2)G_F^2 T^5$$

$$H \approx [(8/3)\pi G\rho_\gamma]^{1/2}$$

$$\rho_\gamma = (\pi^2/30)g_*T^4$$


$$T_f \approx 1 \text{ MeV}$$

$$n/p = \exp\{-\Delta m/T_f\}$$

$$Y_p \approx 2n/(n + p) = 2(n/p)/(n/p + 1)$$

$$\Delta\tau_n = 1\% \rightarrow \Delta Y = 0.75\% (\pm 0.61\%)$$

$$\Delta\tau_n = 1\% \rightarrow \Delta\eta = 17\% (\pm 3.3\%)$$



A.Zalewska, SPC, 17.10.2006


New $\tau_n = (878.5 \pm 0.8)$ s confirms n_b/n_γ from CMB.

Future of Kaon physics

- $K_L^0 \rightarrow \pi^0 \nu \nu$ KEK E391=> JPARC
 - JPARC-P14
- $K^+ \rightarrow \pi^+ \nu \nu$
 - CERN-SPSC-P-326 (a.k.a. NA48/3)
 - 65 signal, 9 ± 3 Bgds/year
 - JPARC-P09
- Transverse μ polarisation in $K^+ \rightarrow \pi^0 \mu^+ \nu$ (T-Violation)
 - JPARC-P06
- Other Initiatives:
 - DANAE (Frascati)
 - OKA (Protvino)

$K^0 \rightarrow \pi^0 \nu \nu$

M.Doroshenko




Spectroscopy - examples

- ➔ Light quark states regularities
- ➔ Heavy charmonium-like states

Striking regularities







- At high M^2 and J :
- Trajectories are linear in (M^2, J) plane
- Trajectories are linear in (M^2, n) plane
- Trajectories are parallel
- The states $I^{GJPC}=0^+0^{++}$ are overpopulated
- The states $I^{GJPC}=0^+2^{++}$ are overpopulated
- Some states are suspicious:
 $\pi_2(1880), \eta_2(1870)$ – candidates to hybrids (Anisovich, Bugg, 2004)
- ≈ 30 states are missed

A. Zaitsev 2006





XYZ




Many new charmonium states: 6 above DD threshold
 + 2 below ($\eta_c(2S)$ and h_c) for last 4 years
 Most of heavy charmonium like states are not explained by theory

A.Zalewska, SPC, 17.10.2006

P.Pakhlov


Near or above the open flavour threshold exotic states are expected to appear in the spectrum: hybrids, molecular states, tetraquarks, ...

X(3872)




in $B \rightarrow KX \rightarrow K\pi^+\pi^-J/\psi$
 $M = 3872.0 \pm 0.6 \pm 0.5$ MeV

BELLE 03, Majumder @ ICHEP 06




$M = 3873.4 \pm 1.4$ (stat) MeV

BABAR 05, Lou @ ICHEP 06



in $p\bar{p} \rightarrow X$
 $\rightarrow \pi^+\pi^-J/\psi$
 $M = 3871.8 \pm 3.1$
 ± 3.0 MeV

D0 04



$M = 3871.3 \pm 0.7$
 ± 0.4 MeV

CDF 04

Kreps @ ICHEP 06

A.Vairo

$X(3872)$: summary of properties



A.Vairo

- $\Gamma_X < 2.3 \text{ MeV}$ (BELLE 03)
- Decay modes: $X \rightarrow \pi^+ \pi^- J/\psi$ (discovery mode), $\pi^+ \pi^- \pi^0 J/\psi$ (BELLE 05), $\gamma J/\psi$ (BELLE 05), $D^0 \bar{D}^0 \pi^0$ (BELLE 06).
- The dominant mode is: $\frac{\mathcal{B}(X \rightarrow D^0 \bar{D}^0 \pi^0)}{\mathcal{B}(X \rightarrow \pi^+ \pi^- J/\psi)} = 9.4_{-4.3}^{+3.6}$ BELLE 06,
but the threshold enhancement peaks at $3875.4 \pm 0.7_{-2.0}^{+1.2}$ MeV: is it $X(3872)$?
- $X \rightarrow \gamma J/\psi \Rightarrow C = +$.
- Angular distribution analyses favour the spin parity: $J = 1+$
BELLE 05, CDF 05
- $\frac{\mathcal{B}(X \rightarrow \pi^+ \pi^- \pi^0 J/\psi)}{\mathcal{B}(X \rightarrow \pi^+ \pi^- J/\psi)} = 1.0 \pm 0.4 \pm 0.3$ BELLE 05
 $\Rightarrow X$ is a mixture of $I = 1$ and $I = 0$ states.
- The substantial $I = 1$ component requires that X contains $u\bar{u}/d\bar{d}$ pairs in addition to hidden charm, which thus qualifies it as a four-quark state. Voloshin 06


Heavy ions

- ➔ RHIC - jet quenching
- ➔ J/ψ suppression
- ➔ Phase transition

Suppression of away-side jet



$$\left(\frac{dE}{dx} \right)_0 \approx 13.8 \pm 3.9 \text{ GeV/fm}$$




$$\left(\frac{dE}{dx} \right)_{\text{cold matter}} \approx 0.5 \text{ GeV/fm}$$

$$\tau_0 = 0.2 \text{ fm/c}$$

Initial Density about 30 times of that in a Cold Au Nucleus

J/ψ production from p-A to Pb-Pb collisions


The study of J/ψ production in p-A collisions at 200, 400 and 450 GeV, by NA3, NA38, NA50 and NA51, gives a “J/ψ absorption cross-section in normal nuclear matter” of 4.18 ± 0.35 mb.



In S-U and *peripheral* Pb-Pb collisions, the data points follow this normal nuclear absorption, which scales with L , the length of nuclear matter crossed by the (pre-resonant) J/ψ.

In central Pb-Pb collisions the L scaling is broken and an “anomalous suppression” sets in.

Comparison with other SPS results




The J/ψ suppression patterns are in fair agreement when plotted versus N_{part}

A.Zalewska, SPC, 17.10.2006

Woehri: ICHEP06

QCD Phase Diagram





News from theory

- ➔ Progress in „practical theory“
- ➔ Gravity/gauge correspondence
- ➔ Little Higgs model = a model of composite Higgs – often quoted in Moscow

On theory side:

V. Rubakov

- Progress in “practical theory”
 - Impressive list of NLO results, crucial NNLO calculations
Marchesini
 - Lattice matches chiral limit and perturbative QCD domain.
With dynamical quarks!
Schierholz



• Gravity/gauge correspondence

Marchesini, Schomerus


- Started up as “theoretical theory” conjecture

adS / CFT

String theory in anti-de Sitter
(5+5)-dimensional space-time

$\mathcal{N}=4$ super-Yang-Mills in Minkowski
4-dimensional space-time

Weak coupling \Leftrightarrow Strong coupling




Calculable
strongly coupled
theory in 4d

V. Rubakov

HEP instrumentation

- ➔ Colliders - recent or near future machines
- ➔ I LC detector - particle flow calorimetry
- ➔ Conference on Instrumentation




A.Skrinsky

A.Zalewska, SPC, 17.10.2006

New Novosibirsk e^+e^- collider VEPP-2000

BEPCII: a double-ring collider


the aim is 4 GeV (tot), $1 \cdot 10^{33} \text{cm}^{-2} \text{s}^{-1}$



PAST: The BEPC was constructed (1984-1988) for both high energy physics and synchrotron radiation research. The machine operated successfully for more than 15 years since 1989. The peak luminosity was $1.2 \times 10^{31} \text{cm}^{-2} \text{s}^{-1}$ @ 1.89 GeV. Detailed study of J/Ψ and Ψ' rare decays.

A. Zalewska, SPC 17-10-2006

A.Skrinsky



Crab cavities will be installed and tested with beam in 2006.

$e^+ 4.1 \text{ A}$




The superconducting cavities will be upgraded to absorb more higher-order mode power up to 50 kW.




Final check

The beam pipes and all vacuum comp higher-current-proof design.



Parameters of RF-Shield

A.Skrinsky



SuperKEKB

$\beta_y^* = \sigma_z = 3 \text{ mm}$

$e^- 9.4 \text{ A}$



The state-of-art ARES copper cavities will be upgraded with higher energy storage ratio to support higher current.

$$L = \frac{\gamma_{\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \frac{I_{\pm} \xi_{\pm y}}{\beta_y^*} \left(\frac{R_L}{R_y} \right)$$

will reach $8 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$.

CLIC for several TeV - is it real solution for lepton-antilepton collisions?

I suspect problems with too many beam-strahlung photons in collision, hence:

- too broad spectrum of e^+e^- collisions.**
- lot of photon-photon “high energy” collisions producing lot of hadrons in each bunch-bunch collision;**

Maybe, not much better than pp collisions of somewhat higher energy?!???

Trends in technology *enable* advances in detectors

examples

- **Segmentation**
 - Vertex elements with 20 μm and smaller features
 - Calorimetry employing silicon elements
 - Micro Pattern Gas Detectors (MPGD) applications
- **Speed**
 - Faster electronics, low noise and low power
- **Integration**
 - Microelectronics
 - Mechanical sophistication
- **Materials**
 - Rad-hard, robust, thin, etc.
- **Radiation immunity**
 - Understanding damage mechanisms and annealing
 - design optimization

Challenges for an ILC Detector

- Vertexing $(h \rightarrow b\bar{b}, c\bar{c}, \tau^+ \tau^-)$
 $\sim 1/5$ r_{beampipe} , $\sim 1/30$ to $1/1000$ pixel size (wrt LHC)

$$\sigma_{ip} = 5\mu\text{m} \oplus 10\mu\text{m} / p \sin^{3/2} \theta$$

- Tracking $(e^+ e^- \rightarrow Zh \rightarrow \ell^+ \ell^- X; \text{incl. } h \rightarrow \text{nothing})$
 $\sim 1/6$ material, $\sim 1/10$ resolution (wrt LHC)

$$\sigma(1/p) = 5 \times 10^{-5} / \text{GeV}$$


- Jet energy (quark reconstruction) \rightarrow calorimetry
 $\sim 1/2$ resolution (wrt LHC)

$$\sigma_E / E = 0.3 / \sqrt{E(\text{GeV})}$$




Particle Flow Calorimetry

- Detector designs are being optimized for the application of Particle Flow Algorithms
 - **Challenge: Calorimeters with very fine readout segmentation**
- The **CALICE** collaboration is building several such calorimeter prototypes
 - a) **Silicon – Tungsten** } **Electromagnetic Calorimeter**
Scintillator – Lead
 - b) **Scintillator** } - Steel Hadronic Calorimeter
RPC
GEM
 - c) Scintillator – Steel Tail Catcher/Muon Tracker
- Tests in particle beams at CERN and FNAL

R.-D. Heuer



Si/W Cal. Assists for Tracking



A.Zalewska, SPC, 17.10.2006

A. Bondar

E. von Toerne

International Instrumentation Conference

C11 proposal:

- Discuss with organizers of successful existing instrumentation conferences whether they support the transition to a truly international conference. Clearly a better solution than creating a completely new conference.
- 2-year cycle of International Instrumentation Conference
- The intermediate year can be used for existing regional conferences.

Conclusion


The LHC start-up should be a highlight of I CHEP08

Medium Response

$$W_{\mu\nu}(q) = \frac{1}{4\pi} \int d^4x e^{iq \cdot x} \langle A | j_\mu^{em}(0) j_\nu^{em}(x) | A \rangle$$

$$F_1(x_B) \quad x_B = -\frac{q^2}{2p \cdot q}$$

Dynamic System:



- **EM emission: Medium response to EM interaction**
 γ production, J/Ψ suppression
- **Hard probes: Medium response to strong interaction**
Jet quenching
- **Soft hadrons: Bulk properties of medium, collective behavior**