

The LAGUNA project (Large Apparatus studying Grand Unification and Neutrino Astrophysics)

Agnieszka Zalewska - IFJ PAN, Poland

Sieć Astrofizyczna, Przegorzały, 15.10.2007

What is LAGUNA

Detector concepts

Research program

Localization of the future large European laboratory

What happens outside Europe?

What is LAGUNA?

The European project „Large Apparatus studying Grand Unification and Neutrino Astrophysics” aiming at defining and realizing this research programme in Europe.


It includes the majority of European groups interested in the construction of the very massive detector ($10^5 - 10^6$ tons) realized in one of the three technologies using liquids: water, liquid argon and liquid scintillator.

No one of the existing European underground laboratories is able to host such a huge detector → a new large underground infrastructure is needed.

The group applied for the RI Design Study in the framework of FP7 (2.05.2007) with the main goal to study possible localizations of the future laboratory together with further R&D for the proposed detector technologies.

The ApPEC roadmap, January 2007

Field/ Experiments	Cost scale (M€)	Desirable start of construction	Remarks
Dark Matter Search: Low background experiments with 1-ton mass	60-100 M€	2011-2013	2 experiments (different nuclei, different techniques), e.g. 1 bolometric, 1 noble liquid; more than 2 worldwide.
Proton decay and low energy neutrino astronomy: Large infrastructure for p- decay and ν astronomy on the 100kt-1Mton scale	400-800 M€	2011-2013	<ul style="list-style-type: none"> - multi-purpose - 3 different techniques; large synergy between them. - needs huge new excavation - expenditures likely also after 2015 <ul style="list-style-type: none"> - worldwide sharing - possibly also accelerator neutrinos in long baseline experiments
The high energy universe: <u>Gamma rays:</u> Cherenkov Telescope Array CTA	100 M€ (South) 50 M€ (North)	first site in 2010	Physics potential well defined by rich physics from present gamma experiments
<u>Charged Cosmic Rays:</u> Auger North	85 M€	2009	Confirmation of physics potential from Auger South results expected in 2007
<u>Neutrinos:</u> KM3NeT	300 M€	2011	FP6 design study. Confirmation of physics potential from IceCube and gamma ray telescopes expected in 2008-2010
Gravitational Waves: Third generation interferometer	250-300 M€	Civil engineering 2012	Conceived as underground laboratory



COLLABORATIVE PROJECT

2.05.2007

Design Study

FP7-INFRASTRUCTURES-2007-1

Proposal title (max 200 characters)

Design of a pan-European
Infrastructure for Large Apparatus
studying Grand Unification and
Neutrino Astrophysics

Proposal acronym

LAGUNA

Type of funding scheme

RI design study implemented as
Collaborative Project

Work programme topics addressed

Deep underground science, particle
physics, astroparticle physics

Name of the coordinating person

Prof. André Rubbia

List of participants:

Participant no.	Participant organisation name	Country
1. ETH Zurich	Swiss Federal Institute of Technology Zurich	Switzerland
2. U-Bern	University of Bern	Switzerland
3. U-Jyväskylä	University of Jyväskylä	Finland
4. U-Oulu	University of Oulu	Finland
5. Rockplan	Kalliosuunnittelu Oy Rockplan Ltd	Finland
6. CEA/ DSM/ DAPNIA	Commissariat à l'Energie Atomique /Direction des Sciences de la Matière	France
7. IN2P3	Institut National de Physique Nucléaire et de Physique des Particules (CNRS/IN2P3)	France
8. MPG	Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V.	Germany
9. TUM	Technische Universität München	Germany
10. U-Hamburg	Universität Hamburg	Germany
11. IFJ PAN	H.Niewodniczanski Institute of Nuclear Physics of the Polish Academy of Sciences, Krakow	Poland
12. IPJ	A.Soltan Institute for Nuclear Studies	Poland
13. US	University of Silesia	Poland
14. UW	Wroclaw University	Poland
15. KGHM CUPRUM	KGHM CUPRUM Ltd Research and Development Centre	Poland
16. IGSMiE PAN	Mineral and Energy Economy Research Institute of the Polish Academy of Sciences	Poland
17. LSC	Laboratorio Subteraneo de Canfranc	Spain
18. UGR	University of Granada	Spain
19. UDUR	University of Durham	United Kingdom
20. U-Sheffield	The University of Sheffield	United Kingdom
21. Technodyne	Technodyne International Ltd	United Kingdom
22. ETL	Electron Tubes	United Kingdom
23. U-Aarhus	University of Aarhus	Denmark
24. AGT	AGT Ingegneria Srl, Perugia	Italy


Przeorganizacja.

Work package no.	Work package title	Type of activity	Lead participant no.	Person-months	Start month	End month
WP1	Management, coordination and assessment	MGT	ETHZ	52	1	36
WP2	Underground Infrastructures and Engineering	RTD	U-Oulu	221	1	35
WP3	Tank Infrastructure and Liquid Handling	RTD	TUM	249	1	35
WP4	Tank Instrumentation and Data Handling	RTD	IN2P3	439	1	35
WP5	Safety and environmental issues	RTD	U-Sheffield	65	1	35
WP6	Science Impact and Outreach	RTD	IFJ PAN	454	1	35
	TOTAL			1480		


Detector concepts

Three liquids: water (MEMPHYS), scintillator (LENA), liquid argon (GLACIER)


MEMPHYS:
Water Cherenkov,
(420 kton - 1 Mton)



LENA:
Liquid Scintillator
(30-70 kton)



GLACIER: Liquid Argon (50 -100 kton)




MEMPHYS - water Cherenkov detector

Concept: initial work for the Frejus laboratory, the SuperKamiokande detector as a prototype, rescaling by a factor up to 20

Advantages: the cheapest target material, mature technology, possible extrapolation to the 1 Mton mass

Challenges: better and cheaper photomultipliers, dopping with $GdCl_3$

Construction: 3-5 tanks, each one with a diameter and a height of 65 m, fiducial mass of 147 ktons read out by 81000 photomultipliers (12" - 30% surface coverage, 20" - 40% coverage)




LENA - the liquid scintillator detector

Concept: initial work for the Pyhäsalmi mine in Finland (underwater placement near Pylos was also considered), the Borexino, Chooz and KamLAND detectors as prototypes, rescaling by a factor 40-50

Advantages: very low energy threshold, good energy resolution, known technology

Challenges: scintillator cleaning, better and cheaper light detection (photomultipliers, light concentrators)

Construction: cylindrical tank 100m long and with a diameter of 30m, fiducial mass of about 50 ktons, readout by 12 000 photomultipliers (20" - 30% surface coverage, with added light concentrates - 50% coverage)



GLACIER - the Liquid Argon detector


Concept: initially developed for Sieroszowice and Gran Sasso, prototype - the ICARUS detector, rescaling by a factor 150

Advantages: very good positional and energetic resolutions
→ imaging topologies, identification of low energy hadrons

Challenges: 20-m long drift of electrons, huge cryogenic installation, dewar thermal insulation

Construction: cylinder
70m in diameter and 20 m
high, total mass - 100 ktons
of Liquid Argon, read out of
the electron ionisation and
light signals (scintillations -
1000 8" PMT, Cherenkov
light - 27000 8" PMT)


Przegorzały.15.10.2007



Research programme

1. Search for the proton decay
2. Studies of the low energy neutrinos from astrophysical sources (SN explosion, Sun, atmospheric neutrinos, relic SN neutrinos in our galaxy) and of the geo-neutrinos
3. Studies of the neutrino properties based on accelerator neutrino beams


Proton decay



MEMPHYS (10 Mtonxyr)



GLACIER (1000 ktonxyr)

LENA (500 ktonxyr)



$p \rightarrow \nu K^+, K^+ \rightarrow \pi^+ \pi^0$ search (SK-I)

typical $p \rightarrow \nu K^+, K^+ \rightarrow \pi^+ \pi^0$ MC event




selection criteria

- 2 e-like ring
- 1 Michel electron
- $85 < m_{\pi^0} < 185 \text{ MeV}/c^2$
- $175 < p_{\pi^0} < 250 \text{ MeV}/c$
- $40 < Q_{\pi^+} < 100 \text{ PE}, Q_{\text{res}} < 70 \text{ PE}$


PROTON DECAY EVENT SIGNATURE

LENA - scintillator






Kaon decay after 18ns

Challenge:
short decay time of
the Kaon (12.8ns)







Kaon decay after 5ns

Proton decay in Liquid Argon



An example of
real event:



Neutrinos from Supernova explosions

1. Supernova physics:

- Gravitational collapse mechanism
- Supernova evolution in time
- Burst detection
- Cooling of the proto-neutron star
- Shock wave propagation
- Black hole formation?


2. Neutrino properties

- Neutrino mass (time of flight delay)
- Oscillation parameters (flavor transformation in SN core and/or in Earth): Type of mass hierarchy and θ_{13} mixing angle

3. Early alert for astronomers

- Pointing to the supernova

Przeorganizacja 15.10.2007




Confirmed
baseline
model...
but still
many
questions



A. Rubbia

Geo-neutrinos

- Antineutrinos from ^{238}U , ^{232}Th i ^{40}K decays inside Earth allow the estimation of the heat generation due to these decays.
- KamLAND experiment provided the first measurement of the flux of geo-neutrinos from the U and Th decays (geo-neutrinos from K decays have energies below the detection threshold in scintillator)



The KamLAND limit for the heat production due to the radioactive decays inside earth < 60 TW

T.Araki et al.,
Nature 436 (2005) 467

KamLAND:


signal 25^{+19}_{-18} , background 127 ± 13

LENA:

expected signal 1000,
background 240(events/year)

Neutrinos from β beam – MEMPHYS

- Acceleration of ${}^6\text{He}$ nuclei (source of antineutrinos) and of ${}^{18}\text{Ne}$ nuclei (source of neutrinos), R&D in the framework of EURISOL DS. (FP6)
- ...But a small obstacle (worth ~1 billion CHF) - the programme requires a serious intervention into the CERN accelerator chain, also problems with poor knowledge of low energy neutrino cross-sections




Low energy neutrinos - cont.

Atmospheric neutrinos - a very big range of E/L,


Search for WIMPS in the SUN and Earth cores

Neutrino astronomy of the Sun


Energy spectrum for the $\nu_e e$ elastic scattering



2006 - spectrum in KamLAND




Possible localizations of the future large underground laboratory



Bedrock zones in the Earth

- Red: very old bedrock, hard crystalline rock: usually very good
- Green: mobile belts (mountains etc), hard rock: fair/variable
- White: sedimentary covers (soft rock): often bad
- Local variations within each zone






Nuclear reactor background

- Relevant mostly for LENA
- Reactor fluxes estimated globally
- Marine reactors irrelevant?


Location	ν (10^8 1/m ² s)
Pyhäsalmi	40
Gran Sasso	54
Frejus	175
Canfranc	196
Boulby	190
Kamioka	408
Sudbury	100
Soudan	33
Pylos	12

Reactor electron anti-neutrino flux density

Prediction for 2015




2005



Przelegni:

Example: baselines from CERN




SUNLAB (Sieroszowice Underground Laboratory)?

Copper, silver and ... salt



Existing big chambers:
Volume $100 \times 15 \times 15 \text{ m}^3$
depth 950 m below the
surface, temp. 35°C ,
humidity $\sim 20\%$




Przekrój geologiczny poprzeczny


1. Czerwony spągowiec; 2. formacja cechsztyńska; 3. pokład soli kamiennej najstarszej (Na 1); 4. trias; 5. trzeciorzęd; 6. czwartorzęd; 7. przypuszczalne dyslokacje uskokuwe

Rock stability - basic question!


Two preliminary geo-mechanical simulations:
W.Pytel (Wrocław), J.Ślizowski, K.Urbańczyk (Kraków)




Effort coefficient distribution (after 30 years)
Rozkład współczynników wyęźnienia (po 30 latach)
model 2/700




Map 51
Criterion 1



Map 52
Criterion 2



Map 53
Criterion 3



Map 54
Criterion 4

Walls movement after 30 years
- by 1.5 m, instant movement
after excavation - 0.145 m

Localization of the future laboratory

Very preliminary sites vs experiments

		Mt Water Cerenkov	50 kt Liquid Scintillator	100 kt Liquid Argon
Fréjus	Tunnel / hard rock	√√√	√√	√√
Gran Sasso	Tunnel / soft rock	√	√√	√
Canfranc	Tunnel	?	?	?
Pyhäsalmi	Mine / hard rock	√	√√√	√√
Boulby	Mine / salt (potash)	?	?	?
Polkowice - Sieroszowice	Mine / salt & rock	√	√√	√√√
Green fields	Own shaft / Hard rock	√	√	√√√

√√√ primary interest; √√ probably; √ unlikely; ? unknown


Outside Europe: Japan - T2K phase II (?)

Accelerator: 4 MW

Detector HiperKamiokande
(1Mton water Cherenkov)

10^6 events
(ν_μ i anti- ν_μ together)

CP
Violation



Neutrino oscillations using the superbeam are an important part of the programme

Outside Europe: USA - DUSEL

DUSEL - Deep Underground Science and Engineering Laboratory

Very rich interdisciplinary programme - from fundamental physics, through biology and engineering studies to the education and outreach.

Six proposed localizations (Homestake, Henderson mine, Soudan, Cascades, San Jacinto, Kimballton), decision soon, startup in 2010 according to the most optimistic scenario.

